



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

III. $\therefore 11\frac{1}{3}\%$ = rate of interest paid.

Solved with varying results by *M. A. Gruber, G. B. M. Zerr, J. E. Baldwin, H. C. Whitaker, H. W. Draughon, I. L. Beverage, and W. F. Bradbury*. Some of the contributors used compound interest.

14. Proposed by Professor G. B. M. ZERR, A. M., Principal of High School, Staunton, Virginia.

A bank by discounting a note of 7% receives for its money a discount equivalent to $7\frac{3}{4}\%$ interest. How long must the note have been discounted before it was due?

Solution by the Proposer.

$7\frac{3}{4}\% - 7\% = \frac{3}{4}\%$. $7\frac{3}{4} : \frac{3}{4} = \$1.00 : \$\frac{3}{8}$, the interest of \$1.00 for the required time at 7%.

$\$1\frac{1}{8} : \$\frac{3}{8} = 12 \text{ months} : 16\frac{2}{3}$ months.

\therefore Time = $16\frac{2}{3}$ months = 1 year 4 month $17\frac{1}{3}$ days.

Also solved with different results by *H. C. Whitaker and P. S. Berg*.

15. Proposed by O. S. KIBLER, Superintendent of Schools, West Middleburg, Logan County, Ohio.

Supposing the town *A* to be 30 mi. from *B*, *B* 25 mi. from *C*, *C* 20 mi. from *A*, where must a house be erected to be equally distant from each of the towns?

Solution by W. A. GRUBER, War Department, Washington, D. C.

The radius of the circumscribed circle of the triangle formed by drawing *AB*, *BC*, and *AC*, is the *distance* required, and the center of this circle is the *place* for the erection of the house.

From Geometry or Trigonometry, we get

$$R = \frac{abc}{4\sqrt{s(s-a)(s-b)(s-c)}},$$

in which *R* represents radius of circumscribed circle of a triangle in terms of the sides of triangle.

Substituting for *a*, *b*, *c*, and *s* [$=\frac{1}{2}(a+b+c)$], the respective values 25, 20, 30, and $37\frac{1}{2}$, we have

$$R = \frac{25 \times 20 \times 30}{4\sqrt{37\frac{1}{2} \times 12\frac{1}{2} \times 17\frac{1}{2} \times 7\frac{1}{2}}}, \text{ which reduced, becomes}$$

$$R = \frac{40}{\sqrt{7}} = \frac{40\sqrt{7}}{7} = 15.11857 \text{ mi.}$$

Also solved by *H. C. Whitaker, G. B. M. Zerr, Seth Pratt, J. F. W. Scheffer, J. W. Watson, and P. S. Berg*.

PROBLEMS.

22. Proposed by E. S. Loomis, A. M., Ph.D., Professor of Mathematics, Baldwin University, Berea, Ohio.

A borrows \$1000 from *B* for 10 years, on which he pays 4% semi-annually.

A immediately loans the \$1000 to *C* for 10 years, who agrees to pay to *A* \$12 $\frac{1}{2}$ on the first of each month for 120 mos. or 10 yrs., at which time the whole debt is considered canceled, *C* no longer being, in any way, indebted to *A*. Upon the receipt of each of the \$12 $\frac{1}{2}$ payments made by *C*, *A* immediately relays it to *D*, *E*, *F*, etc., upon the same conditions as he loaned the \$1000 to *C*; at the end of 120 mos. all who are indebted to *A* pay up in full all due him, and he (*A*) pays *B* the principal, all interest having been paid when due.

Query: How many dollars has he in hand?

23. Proposed by H. C. Whitaker, Professor of Mathematics, Manual Training School, Philadelphia, Pennsylvania.

A rectangular hall 80 feet long, 40 feet wide and 12 feet high has a spider in one corner of the ceiling. How long will it take the spider to crawl to the opposite corner on the floor if he crawls a foot in a second on the wall and two feet in a second on the floor?

24. Proposed by Mrs. Mary E. Hogsett, Danville, Kentucky.

On January 4, 1889, it was noticed that a clock was 15 minutes fast. On March 1, 1894, it was found to be six and one half minutes slow. When and what time was accurate time?

Solutions to these problems should be received on or before July 1st.

ALGEBRA.

Conducted by J. M. OOLAW, Monterey, Va. All contributions to this department should be sent to him.

SOLUTIONS TO PROBLEMS.

11. Proposed by ISAAC L. BEVERAGE, Monterey, Virginia.

Two men, A and B , had a money-box, containing \$210, from which each drew a certain sum daily; this sum being fixed for each, but different for the two. After six weeks, the box was empty. Find the sum which each man drew daily from the box, knowing that A alone would have emptied it five weeks earlier than B alone,

- I. Solution by W. L. HARVEY, Portland, Maine.

Let x = amount B drew out daily, mx = amount A drew out daily.

$$\text{Then, } \frac{210}{x} - \frac{210}{mx} = 35 \dots (1), \text{ and } \frac{210}{x(m+1)} = 42 \dots (2).$$

Solving $m = 1\frac{1}{2}$, whence $x = \$2$, and $mx = \$3$.

- II. Solution by P. S. BERG, Apple Creek, Ohio.

Let x = what A drew out daily, y = what B drew out daily.

$$\text{Then, } 42x + 42y = \$210 \dots (1), \text{ and } \frac{210}{y} - \frac{210}{x} = 35 \dots (2).$$

Whence $x = 3$, and $y = 2$. $\therefore A$ drew out daily \$3, and B \$2.

Also solved by M. A. GRUBER, H. W. DRAUGHON, H. C. WHITAKER, C. E. MYERS, G. B. M. ZERR, A. L. FOOTE, and ROBERT J. ALEY.

12. Proposed by F. M. SHIELDS, Coopwood, Mississippi.

Three lads, A , B , and C , each climbed to the top of an upright pole: A 's pole was 20 feet high, B 's 60 feet, and C 's pole was 100 feet high. They all started at the same time, and each climbed up a part of the way, at the same rate of speed per minute, and after each rested five minutes, they ascended to the tops of their respective poles, at the same rate of speed per minute, when they found that each had consumed